



# TAKHZYRO™

(lanadelumab-flyo) injection

## NOW APPROVED

### TAKHZYRO IS A FIRST-OF-ITS-KIND MAB PREVENTIVE TREATMENT FOR HEREDITARY ANGIOEDEMA (HAE)<sup>1</sup>

#### JOIN US TO LEARN MORE ABOUT:

- The pathophysiology of hereditary angioedema (HAE) and the impact it has on patients
- TAKHZYRO (lanadelumab-flyo): A first-of-its-kind mAb injectable medicine for prophylaxis to prevent attacks of HAE in patients 12 years and older
  - The recommended starting dose is 300 mg every 2 weeks. A dosing interval of 300 mg every 4 weeks is also effective and may be considered if the patient is well-controlled (e.g., attack free) for more than 6 months.

TO REGISTER, PLEASE CALL  
877-761-7925 OR VISIT  
[www.programsvp.com/2695-39](http://www.programsvp.com/2695-39)

TAKHZYRO is available as a 300 mg dose

*This program is open to healthcare professionals only. **Please be advised that spouses and other guests are not permitted to participate.** Thank you in advance, and we look forward to your participation.*

**This is not a CME program.**

#### INDICATION

TAKHZYRO (lanadelumab-flyo) is indicated for prophylaxis to prevent attacks of hereditary angioedema (HAE) in patients  $\geq 12$  years of age.

#### IMPORTANT SAFETY INFORMATION

**Hypersensitivity reactions** have been observed. In case of a severe hypersensitivity reaction, discontinue TAKHZYRO administration and institute appropriate treatment.

**Please see next page for additional Important Safety Information.**



#### PRESENTER:

William R. Lumry, MD  
Founder  
Allergy & Asthma Specialists of Dallas  
Dallas, TX



#### WHEN:

Friday, April 26, 2019  
7:00 PM ET



#### WHERE:

Washington Duke Inn & Golf Club  
3001 Cameron Boulevard  
Durham, NC 27705



TAKHZYRO™  
(lanadelumab-flyo) injection

## IMPORTANT SAFETY INFORMATION (cont'd)

**Adverse Reactions:** The most commonly observed adverse reactions ( $\geq 10\%$  and higher than placebo) associated with TAKHZYRO were injection site reactions consisting mainly of pain, erythema, and bruising at the injection site; upper respiratory infection; headache; rash; myalgia; dizziness; and diarrhea. Less common adverse reactions observed included elevated levels of transaminases; one patient discontinued the trial for elevated transaminases.

**Use in Specific Populations:** The safety and efficacy of TAKHZYRO in pediatric patients <12 years of age have not been established.

No data are available on TAKHZYRO in pregnant women. No data are available on the presence of lanadelumab in human milk or its effects on breastfed infants or milk production.

To report SUSPECTED ADVERSE REACTIONS, contact Dyax Corp. (a wholly-owned, indirect subsidiary of Shire plc) at 1-800-828-2088, or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).

Please see full [Prescribing Information](#), including information for patients.

**Reference: 1.** TAKHZYRO (lanadelumab-flyo) injection [prescribing information].  
Lexington, MA: Shire Pharmaceuticals, Inc; 2018.

TAKHZYRO<sup>™</sup>  
(lanadelumab-flyo) injection

Program sponsored by

**Shire**

©2018 Shire US Inc., 300 Shire Way, Lexington, MA 02421.  
1-800-828-2088. All rights reserved. TAKHZYRO is a trademark or registered trademark of Dyax Corp., a wholly-owned, indirect subsidiary of Shire plc. SHIRE, and the associated logos are trademarks or registered trademarks of Shire.

S43349 10/18